

THEOREM 3 Mean Value Theorem for Derivatives

If $y = f(x)$ is continuous at every point of the closed interval $[a, b]$ and differentiable at every point of its interior (a, b) , then there is at least one point c in (a, b) at which

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

EXAMPLE 1 Exploring the Mean Value Theorem

Show that the function $f(x) = x^2$ satisfies the hypotheses of the Mean Value Theorem on the interval $[0, 2]$. Then find a solution c to the equation

on this interval.

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$a = 0$$

$$b = 2$$

$$f'(c) = 2$$

$$f'(x) = 2x$$

$$2x = 2$$

$$x = 1$$

$$c = 1$$

EXAMPLE 2 Exploring the Mean Value Theorem

Explain why each of the following functions fails to satisfy the conditions of the Mean Value Theorem on the interval $[-1, 1]$.

(a) $f(x) = \sqrt{x^2} + 1$

$|x|$
corner

(b) $f(x) = \begin{cases} x^3 + 3 & \text{for } x < 1 \\ x^2 + 1 & \text{for } x \geq 1 \end{cases}$

not continuous

EXAMPLE 3 Applying the Mean Value Theorem

Let $f(x) = \sqrt{1 - x^2}$, $A = (-1, f(-1))$, and $B = (1, f(1))$. Find a tangent to f in the interval $(-1, 1)$ that is parallel to the secant AB .

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$f'(c) = \frac{0 - 0}{2} = 0$$

$$f'(x) = \frac{-x}{\sqrt{1 - x^2}}$$

$$x = 0 \quad f(0) = 1$$

y = 1

Physical Interpretation

If we think of the difference quotient $(f(b) - f(a))/(b - a)$ as the average change in f over $[a, b]$ and $f'(c)$ as an instantaneous change, then the Mean Value Theorem says that the instantaneous change at some interior point must equal the average change over the entire interval.

EXAMPLE 4 Interpreting the Mean Value Theorem

If a car accelerating from zero takes 8 sec to go 352 ft, its average velocity for the 8-sec interval is $352/8 = 44$ ft/sec, or 30 mph. At some point during the acceleration, the theorem says, the speedometer must read exactly 30 mph (Figure 4.15).

DEFINITIONS Increasing Function, Decreasing Function

Let f be a function defined on an interval I and let x_1 and x_2 be any two points in I .

1. f **increases** on I if $x_1 < x_2 \Rightarrow f(x_1) < f(x_2)$.
2. f **decreases** on I if $x_1 < x_2 \Rightarrow f(x_1) > f(x_2)$.

COROLLARY 1 Increasing and Decreasing Functions

Let f be continuous on $[a, b]$ and differentiable on (a, b) .

1. If $f' > 0$ at each point of (a, b) , then f increases on $[a, b]$.
2. If $f' < 0$ at each point of (a, b) , then f decreases on $[a, b]$.

EXAMPLE 5 Determining Where Graphs Rise or Fall

The function $y = x^2$ (Figure 4.16) is

EXAMPLE 6 Determining Where Graphs Rise or Fall

Where is the function $f(x) = x^3 - 4x$ increasing and where is it decreasing?

COROLLARY 2 Functions with $f' = 0$ are Constant

If $f'(x) = 0$ at each point of an interval I , then there is a constant C for which $f(x) = C$ for all x in I .

COROLLARY 3 Functions with the Same Derivative Differ by a Constant

If $f'(x) = g'(x)$ at each point of an interval I , then there is a constant C such that $f(x) = g(x) + C$ for all x in I .

EXAMPLE 7 Applying Corollary 3

Find the function $f(x)$ whose derivative is $\sin x$ and whose graph passes through the point $(0, 2)$.

EXAMPLE 7 Applying Corollary 3

Find the function $f(x)$ whose derivative is $\sin x$ and whose graph passes through the point $(0, 2)$.

DEFINITION Antiderivative

A function $F(x)$ is an **antiderivative** of a function $f(x)$ if $F'(x) = f(x)$ for all x in the domain of f . The process of finding an antiderivative is **antidifferentiation**.

EXAMPLE 8 Finding Velocity and Position

Find the velocity and position functions of a body falling freely from a height of 0 meters under each of the following sets of conditions:

- (a) The acceleration is 9.8 m/sec^2 and the body falls from rest.
- (b) The acceleration is 9.8 m/sec^2 and the body is propelled downward with an initial velocity of 1 m/sec .